

CHORIOAMNIONITIS, MECHANICAL VENTILATION AND SEPSIS AS MODULATORS OF BRONCHOPULMONARY DYSPLASIA

Fernandes G, Firmino, I, Margotto, P.

HOSPITAL REGIONAL DA ASA SUL (HRAS), BRASILIA, BRAZIL. | HOSPITAL MATERNO-INFANTIL DE BRASÍLIA

Table 1. Obstetric and neonatal characteristics of the study population according to the presence of histologic chorioamnionitis

Variables	With chorioamnionitis (n 23)	Without chorioamnionitis (n 28)	P
Maternal age	27,5 ±8,1	25,6 ±6,9	0,48
Pregnancies	3,0 ±1,8	1,8 ±1,5	0,04
Prenatal corticosteroids	14 (60,9%)	14 (50%)	0,43
Cesarean	6 (26,1%)	19 (67,9%)	0,003
Resuscitation	13 (56,5%)	18 (64,3%)	0,57
Gestational age	27,9 ±2	29,2 ±1,9	0,02
Birth weight (g)	1059,1 ±361,4	1168,6 ±329,7	0,27
Height	36,4 ±3,9	39,7 ±5,5	0,10
Head circumference	25,5 ±2,4	27,0 ±2,4	0,03
PIG	3 (13%)	3 (10,7%)	0,79
Male	10 (43,5%)	13 (46,4%)	0,6
Days in NICU	67,14 ±37,7	63,1 ±42,0	0,84
FiO2 (%)	61,3 ±24,3	54,0 ±25,4	0,30
Days with O2	32,13 ±37,4	22,9 ±29,0	0,33

Table 2. Clinical outcome in the neonatal period

Variable	With chorioamnionitis (n 23)	Without chorioamnionitis (n28)	Relative risk (IC* 95%)	P
Surfactant	20 (87%)	20 (71,4%)	2,5 (0,61 - 11,5)	0,18
RDS [‡]	20 (87%)	23 (82,1%)	1,44 (0,3 - 6,8)	0,6
MV (days) [§]	11,5 ±16,1	10,1 ±17,5	-	0,76
CPAP (days) ^{¶¶}	7,57 ±14,6	4,4 ±4,4	-	0,29
O2 in 28 days	8 (34,8%)	8 (29,6%)	1,26 (0,38 - 4,16)	0,69
O2 in 36 weeks	6 (27,3%)	4 (15,4%)	2,0 (0,49 - 8,5)	0,3
Early sepsis	21 (91,3%)	13 (50%)	10,5 (2 - 54,2)	0,002
PDA [‡]	9 (42,9%)	4 (16%)	3,9 (0,99 - 15,5)	0,04
Death	6 (26,1%)	4 (14,3%)	2,1 (0,51 - 8,7)	0,24

PDA - Persistent Ductus Arteriosus. VM – Mechanical ventilation. RDS - Respiratory distress syndrome

Reference

Miralles RE, Hodge R, Kotecha S. Antenatal inflammation and infection in chronic lung disease of prematurity. Child: Care, Health and Development, 28: 11-15, 2002

Dammann O, Leviton A, Gappa M, Dammann CEL. Lung and brain damage in preterm newborns, and their association with gestacional age, prematurity subgroup, infection/inflammation and long term outcome. International Journal of Obstetrics and Gynaecology, 112: 4-9, 2005

Zanardo V, Vedovato S, Cosmi E, Litta P, Callin F, Trevisanuto D, Chiarelli S. Preterm premature rupture of membranes, chorioamnion inflammatory scores and neonatal respiratory outcome. International Journal of Obstetrics and Gynaecology, 117: 94-98, 2010

Prendergast M, May C, Broughton S, Pollina E, Milner AD, Rafferty GF, Greenough. Chorioamnionitis, lung function and bronchopulmonary dysplasia in prematurely born infants. Arch Dis Child Fetal Neonatal 96: F270-274, 2011

Viscardi RM. Perinatal inflammation and lung injury. Seminars in Fetal and Neonatal Medicine, 1-6, 2011

Paul DA, Zook K, Mackley A, Locke RG. Reduced mortality and increased BPD with histological chorioamnionitis and leukocytosis in very-low-birth-weight infants. Journal of Perinatology, 30: 58-62, 2010.

Paanannem R, Husa AK, Vuolteenaho R, Herva R, Kaukola T. Blood Cytokines during the Perinatal Period in Very Preterm Infants: Relationship of Inflammatory Response and Bronchopulmonary Dysplasia. The Journal of Pediatrics, 154: 39 - 43, 2009.

Kramer BW, Kallapur S, Newnham J, Jobe AH. Prenatal inflammation and lung development. Seminars in Fetal and Neonatal Medicine, 14: 2-7, 2009.

Kramer BW. Antenatal inflammation and lung injury: prenatal origin of neonatal disease. Journal of Perinatology, 28: S21-S27, 2008.

Method

An observational retrospective cohort of neonates with 26-32 weeks, born in Hospital Materno Infantil de Brasilia, Brazil, from April to October 2011. Statistical analysis was performed using SPSS version 16.0 software, and used the chi-square tests of association, t-Sudent, the hazard ratio with confidence interval. The significance level was 0.05.

Results and Discussion

Seventy eighth mothers conceived 86 infants with gestational age between 26 and 32 weeks from April to October 2011. Only 51 patients had the histopathological results in your medical records and 23 mothers had the histological chorioamnionitis. We studied 51 preterm infants, and 23 underwent antenatal infection. Mothers with chorioamnionitis had a higher number of previous pregnancies were less submitted to cesarean section. Receiving corticosteroids as were mothers without infection. Newborns who underwent maternal infection, were born with gestational age and lower weight but a smaller number needed resuscitation at birth. Had 10 times more early-onset sepsis, a higher incidence of patent ductus arteriosus and a 2 times greater tendency to develop bronchopulmonary dysplasia.The association bettween chorioamnionitis end Persistent Ductus

Arteriosus has been observed in other studies and can be explained by the increase in prostaglandin levels and tumor necrosis factor in the fetal circulation. It is also known that there is an association between persistent ductus arteriosus and bronchopulmonary dysplasia. The left-right shunt at the ductus arteriosus increases blood flow in the lung, with subsequent reduction in lung compliance and increased resistance, thereby raising the need for mechanical ventilation and supplemental oxygen, further aggravating lung injury.

Conclusion

The premature newborns have several risk factors for the development of lung injury. The chorioamnionitis is clearly a contributing factor, despite the difficulty of finding an isolated risk factor in the pathogenesis of bronchopulmonary dysplasia. The evaluation studies to help clarify the pathogenesis of these diseases is essential to improve neonatal interventions. It has a major impact on the increased survival of premature infants, the quality children’s life and consequently bring positive impact on adult health. Despite the small sample limit our findings, it opens the way for further research, in order to have health practices increasingly based on scientific evidence.

Richardson BS. Preterm histologic chorioamnionitis: impact on cord gas and pH values and neonatal outcome. American Journal Obstetrics and Gynaecology, 195: 1357 - 65, 2006

Gordon A, Lahra M, Greenow CR, Jeffery H. Histological Chorioamnionitis is increased at extremes of gestation in stillbirth: a population based study. Infect Dis Obstet Gynecol. 2011;456728, 2011

Villar J. Maternal and Neonatal individuals risks and benefits associated with caesarean delivery: multicentre prospective study. BMJ, 335: 1025 - 27, 2007.

Jobe AH, Kallapur SG. Contribution of inflammation to lung injury and development. Arch Dis Child Fetal Neonatal, 91: F132-F135, 2006

Vigneswaran R. Infection and preterm birth: evidence of a common causal relationship with bronchopulmonary dysplasia and cerebral palsy. Child health, 36: 293-296, 2000

Bancalari E. Displasia broncopulmonar: um velho problema de cara nova. Jornal de Pediatria, 82: 2-3, 2006

May C, Patel S, Kennedy C, Pollina E, Rafferty G, Paacock JL, Greenough A. Prediction of bronchopulmonary dysplasia. Arch Dis Child Fetal Neonatal, 96: F410 -F416, 2011

Jobe AH. Antenatal associations with lung maturation and infection. Journal of Perinatology, 25: S31-S35, 2005

Lee HJ, Kim EK, Kim HS, Choi CW, Kim B, Choi JH. Chorioamnionitis, respiratory syndrome and bronchopulmonary dysplasia in extremely low birth weight infants. Journal of Perinatology, 31: 166-170, 2011

Watterberg KL, Demers LM, Scott SM, Murphy S. Chorioamnionitis and early lung inflammation in infants in whom bronchopulmonary dysplasia develops. Pediatrics 1996 97:210-5,1996

Bancalari E, Claure N, Sosenko IRS. Bronchopulmonary dysplasia: changes in pathogenesis, epidemiology and definition. Seminars in Neonatology, 8: 63-71, 2003.